

CONNECTED COMPONENTS OF THE SPACE OF SURFACE GROUP REPRESENTATIONS

NAN-KUO HO AND CHIU-CHU MELISSA LIU

ABSTRACT. Let G be a connected, compact, semisimple Lie group. It is known that for a compact closed orientable surface Σ of genus $l > 1$, the order of the group $H^2(\Sigma, \pi_1(G))$ is equal to the number of connected components of the space $\text{Hom}(\pi_1(\Sigma), G)/G$ which can also be identified with the moduli space of gauge equivalence classes of flat G -bundles over Σ . We show that the same statement for a closed compact nonorientable surface which is homeomorphic to the connected sum of k copies of the real projective plane, where $k \neq 1, 2, 4$, can be easily derived from a result in A. Alekseev, A. Malkin and E. Meinrenken's recent work on Lie group valued moment maps.

1. INTRODUCTION

Let Σ be a closed compact orientable surface of genus $l > 1$. In [Go], W.M. Goldman conjectured that for any connected complex semisimple Lie group G , there is a bijection

$$\pi_0(\text{Hom}(\pi_1(\Sigma), G)) \rightarrow H^2(\Sigma; \pi_1(G)) \cong \pi_1(G).$$

In [Li], J. Li proved the above conjecture by Goldman as well as the following:

Theorem 1 ([Li, Theorem 0.5]). *Let Σ be a closed orientable Riemann surface of genus $l > 1$. Let G be a connected, compact, semisimple Lie group. Then there is a bijection*

$$\pi_0(\text{Hom}(\pi_1(\Sigma), G)/G) \rightarrow H^2(\Sigma; \pi_1(G)) \cong \pi_1(G),$$

where G acts on $\text{Hom}(\pi_1(\Sigma), G)$ by conjugation.

Geometrically, $\text{Hom}(\pi_1(\Sigma), G)/G$ can be identified with the moduli space of gauge equivalence classes of flat G -bundles over Σ , where a flat G -bundle is a principal G -bundle together with a flat connection. It is known that there is a one-to-one correspondence between topological principal G -bundles over Σ and elements in $H^2(\Sigma; \pi_1(G)) \cong \pi_1(G)$. In [Li], J. Li combined the argument in [Ra] and [AB] to prove that the moduli space of gauge equivalence classes of flat connections on a fixed underlying topological principal G -bundle over Σ is nonempty and connected, where Σ and G are as in Theorem 1.

The main result in this paper is the following analogue of Theorem 1.

Date: February 1, 2008.

1991 Mathematics Subject Classification. 53.

Key words and phrases. moduli space of flat G connections.

Theorem 2. *Let Σ be a closed compact nonorientable surface which is homeomorphic to the connected sum of k copies of the real projective plane, where $k \neq 1, 2, 4$. Let G be a connected, compact, semisimple Lie group. Then there is a bijection*

$$\pi_0(\text{Hom}(\pi_1(\Sigma), G)/G) \rightarrow H^2(\Sigma; \pi_1(G)) \cong \pi_1(G)/2\pi_1(G),$$

where $2\pi_1(G)$ denote the subgroup $\{k^2 \mid k \in \pi_1(G)\}$ of the finite abelian group $\pi_1(G)$.

One can see the isomorphism $H^2(\Sigma; \pi_1(G)) \cong \pi_1(G)/2\pi_1(G)$ as follows. By the universal coefficient theorem of cohomology, we have a short exact sequence

$$0 \rightarrow \text{Ext}(H_1(\Sigma; \mathbb{Z}), \pi_1(G)) \rightarrow H^2(\Sigma; \pi_1(G)) \rightarrow \text{Hom}(H_2(\Sigma; \mathbb{Z}), \pi_1(G)) \rightarrow 0,$$

where $H_1(\Sigma, \mathbb{Z}) \cong \mathbb{Z}^{k-1} \oplus \mathbb{Z}/2\mathbb{Z}$, and $H_2(\Sigma, \mathbb{Z}) = 0$. So we have

$$H^2(\Sigma, \pi_1(G)) \cong \text{Ext}(\mathbb{Z}/2\mathbb{Z}, \pi_1(G)) \cong \pi_1(G)/2\pi_1(G).$$

Let Σ be a closed compact nonorientable surface. It is known that there is a one-to-one correspondence between topological principal G -bundles over Σ and elements in $H^2(\Sigma; \pi_1(G)) \cong \pi_1(G)/2\pi_1(G)$. The geometric interpretation of Theorem 2 is that the moduli space of gauge equivalence classes of flat connections on a fixed underlying topological principal G -bundle over Σ is nonempty and connected, where Σ and G are as in Theorem 2. More details of this geometric interpretation will be given in Appendix A.

We will show that both Theorem 1 and Theorem 2 can be easily derived from the following Fact 3, without identifying $\text{Hom}(\pi_1(\Sigma), G)/G$ with the moduli space of gauge equivalence classes of flat G -bundles over Σ .

Fact 3. *Let G be a compact, semisimple, connected and simply connected Lie group. Let l be a positive integer. Then the commutator map $\mu_G^l : G^{2l} \rightarrow G$ defined by*

$$(1) \quad \mu_G^l(a_1, b_1, \dots, a_l, b_l) = a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1}$$

is surjective, and $(\mu_G^l)^{-1}(g)$ is connected for all $g \in G$.

The surjectivity follows from Goto's commutator theorem [HM, Theorem 6.55]. The commutator map μ_G^l in Fact 3 is a group valued moment map of the q-Hamiltonian G -space G^{2l} in the sense of A. Alekseev, A. Malkin, and E. Meinrenken [AMM]. By [AMM, Theorem 7.2], all the fibers of the moment map of a connected q-Hamiltonian G -space are connected if G is a compact, connected and simply connected Lie group.

This paper is organized as follows. In Section 2, we explain how Fact 3 implies Theorem 1 as well as the genus one case. In Section 3 and Section 4, we derive Theorem 2 from Fact 3 for odd k and even k , respectively. In Section 5, we discuss the cases $k = 1, 2, 4$ which are not covered by our main result. In Appendix A, we give algebraic and geometric interpretations of the obstruction map used in our proof.

Acknowledgment: It is a pleasure to thank J.-K. Yu, who motivated and guided this work. We also wish to thank L. Jeffrey and E. Meinrenken for helpful conversations and valuable suggestions.

2. COMPACT ORIENTABLE SURFACES

In this section, Σ is a closed Riemann surface with genus $l > 0$. Note that $\text{Hom}(\pi_1(S^2), G)/G$ consists of a single point, so Theorem 1 is not true for the genus zero case.

Let G be a connected, compact, semisimple Lie group, and let $\rho : \tilde{G} \rightarrow G$ be the universal covering map which is also a group homomorphism. Then $\tilde{G}$ is a compact, semisimple, connected and simply connected Lie group, and $\text{Ker}\rho$ is a subgroup of the center $Z(\tilde{G})$ of $\tilde{G}$ which is abelian.

Let $\mu_G^l : G^{2l} \rightarrow G$ be the commutator map defined as in (1), and let e be the identity element of G . Then $\text{Hom}(\pi_1(\Sigma), G)$ can be identified with

$$(\mu_G^l)^{-1}(e) = \{(a_1, b_1, \dots, a_l, b_l) \in G^{2l} \mid a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1} = e\}.$$

There is an obstruction map (see e.g. [Go])

$$o_2 : \text{Hom}(\pi_1(\Sigma), G) \cong (\mu_G^l)^{-1}(e) \rightarrow \text{Ker}\rho$$

defined by

$$(a_1, b_1, \dots, a_l, b_l) \in (\mu_G^l)^{-1}(e) \mapsto \tilde{a}_1 \tilde{b}_1 \tilde{a}_1^{-1} \tilde{b}_1^{-1} \cdots \tilde{a}_l \tilde{b}_l \tilde{a}_l^{-1} \tilde{b}_l^{-1} \in \text{Ker}\rho,$$

where $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l)$ is a preimage of $(a_1, b_1, \dots, a_l, b_l)$ under $\rho^{2l} : \tilde{G}^{2l} \rightarrow G^{2l}$. It is easily checked that the definition does not depend on the choice of $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l)$ and o_2 descends to a continuous map

$$\bar{o}_2 : \text{Hom}(\pi_1(\Sigma), G)/G \rightarrow \text{Ker}\rho.$$

For each $k \in \text{Ker}\rho$, there is a surjective continuous map

$$(\mu_{\tilde{G}}^l)^{-1}(k) \rightarrow \bar{o}_2^{-1}(k)$$

which is the restriction of $\tilde{G}^{2l} \xrightarrow{\rho^{2l}} G^{2l} \rightarrow G^{2l}/G$, where G acts on G^{2l} by diagonal conjugation. By Fact 3, $(\mu_{\tilde{G}}^l)^{-1}(k)$ is nonempty and connected for each $k \in \text{Ker}\rho$, so $\bar{o}_2$ is surjective, and $\bar{o}_2^{-1}(k)$ is connected for each $k \in \text{Ker}\rho \cong \pi_1(G)$. This gives Theorem 1 as well as the genus one case.

3. CONNECTED SUM OF A ODD NUMBER OF COPIES OF THE REAL PROJECTIVE PLANE

In this section, Σ is a closed compact nonorientable surface which is homeomorphic to the connected sum of $2l+1$ copies of RP^2 , or equivalently, the connected sum of a Riemann surface of genus l and RP^2 , where $l > 0$.

Let G be a connected, compact, semisimple Lie group, and let $\rho : \tilde{G} \rightarrow G$ be the universal covering map. The space $\text{Hom}(\pi_1(\Sigma), G)$ can be identified with

$$X = \{(a_1, b_1, \dots, a_l, b_l, c) \in G^{2l+1} \mid a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1} c^2 = e\},$$

where e is the identity element of G .

Let $K = \text{Ker} \rho \cong \pi_1(G)$, and let $2K$ be the subgroup $\{k^2 \mid k \in K\}$ of the finite abelian group K . There is an obstruction map

$$o_2 : \text{Hom}(\pi_1(\Sigma), G) \cong X \rightarrow K/2K$$

defined by

$$(a_1, b_1, \dots, a_l, b_l, c) \in X \mapsto \tilde{a}_1 \tilde{b}_1 \tilde{a}_1^{-1} \tilde{b}_1^{-1} \cdots \tilde{a}_l \tilde{b}_l \tilde{a}_l^{-1} \tilde{b}_l^{-1} \tilde{c}^2 \in K/2K,$$

where $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l, \tilde{c})$ is a preimage of $(a_1, b_1, \dots, a_l, b_l, c)$ under $\rho^{2l+1} : \tilde{G}^{2l+1} \rightarrow G^{2l+1}$. It is easily checked that the definition does not depend on the choice of $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l, \tilde{c})$ and o_2 descends to a continuous map

$$\bar{o}_2 : \text{Hom}(\pi_1(\Sigma), G)/G \rightarrow K/2K.$$

For each $k \in K$, define

$$\tilde{X}_k = \{(a_1, b_1, \dots, a_l, b_l, c) \in \tilde{G}^{2l+1} \mid a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1} c^2 = k\}.$$

Let $\bar{k}$ denote the image of k under $K \rightarrow K/2K$. Then there is a surjective continuous map

$$\tilde{X}_k \rightarrow \bar{o}_2^{-1}(\bar{k})$$

which is the restriction of $\tilde{G}^{2l+1} \xrightarrow{\rho^{2l+1}} G^{2l+1} \rightarrow G^{2l+1}/G$, where G acts on G^{2l+1} by diagonal conjugation. Let $\tilde{e}$ denote the identity element of $\tilde{G}$. For each $k \in K$, $\tilde{X}_k$ contains $(\mu_{\tilde{G}}^l)^{-1}(k) \times \{\tilde{e}\}$ which is nonempty by Fact 3. Therefore, $\bar{o}_2$ is surjective. It remains to show that $\tilde{X}_k$ is connected for all $k \in K$, which will imply $\bar{o}_2^{-1}(\bar{k})$ is connected for all $\bar{k} \in K/2K$.

Let $Q' : \tilde{G}^{2l+1} \rightarrow \tilde{G}$ be the projection to the last factor. The restriction of Q' to $\tilde{X}_k$ gives a map $Q : \tilde{X}_k \rightarrow \tilde{G}$, $(a_1, b_1, \dots, a_l, b_l, c) \mapsto c$. Fact 3 implies that Q is surjective and $Q^{-1}(c)$ is connected for all $c \in \tilde{G}$. Let $q \in T$ be a square root of $k \in Z(\tilde{G}) \subset T$, where we fix a maximal torus T of $\tilde{G}$. To prove that $\tilde{X}_k$ is connected, it suffices to show that for any $c \in \tilde{G}$, there is a path $\gamma : [0, 1] \rightarrow \tilde{X}_k$ such that $\gamma(0) \in Q^{-1}(q)$ and $\gamma(1) \in Q^{-1}(c)$.

For any $c \in \tilde{G}$, there exists $g \in \tilde{G}$ such that $g^{-1}cg \in T$. Let $\mathfrak{g}$ and $\mathfrak{t}$ be the Lie algebras of $\tilde{G}$ and T , respectively. Let $\exp : \mathfrak{g} \rightarrow \tilde{G}$ be the exponential map. Then $g^{-1}cgq^{-1} = \exp \xi$ for some $\xi \in \mathfrak{t}$. Let W be the Weyl group of $\tilde{G}$, and let $w \in W$ be a Coxeter element. The linear map $w : \mathfrak{t} \rightarrow \mathfrak{t}$ has no eigenvalue equal to 1 [Hu, Section 3.16], so there exists $\xi' \in \mathfrak{t}$ such that $w \cdot \xi' - \xi' = \xi$. Recall that $W = N(T)/T$, where $N(T)$ is the normalizer of T in $\tilde{G}$, so $w = aT \in N(T)/T$ for some $a \in \tilde{G}$. We have

$$a \exp(t\xi') a^{-1} \exp(-t\xi') = \exp(t\xi)$$

for any $t \in \mathbb{R}$.

The group $\tilde{G}$ is connected, so there exists a path $\tilde{g} : [0, 1] \rightarrow \tilde{G}$ such that $\tilde{g}(0) = \tilde{e}$ and $\tilde{g}(1) = g$. Define $\gamma : [0, 1] \rightarrow \tilde{G}^{2l+1}$ by

$$\gamma(t) = (a(t), b(t), \tilde{e}, \dots, \tilde{e}, c(t)),$$

where

$$a(t) = \tilde{g}(t)a\tilde{g}(t)^{-1}, \quad b(t) = \tilde{g}(t)\exp(-2t\xi')\tilde{g}(t)^{-1}, \quad c(t) = \tilde{g}(t)q\exp(t\xi)\tilde{g}(t)^{-1}.$$

Then the image of γ lies in $\tilde{X}_k$, $\gamma(0) = (a, \tilde{e}, \tilde{e}, \dots, \tilde{e}, q) \in Q^{-1}(q)$, and

$$\gamma(1) = (gag^{-1}, g\exp(-2\xi')g^{-1}, \tilde{e}, \dots, \tilde{e}, c) \in Q^{-1}(c).$$

Remark 4. Let M be a manifold, and let G be a compact Lie group. By [AMM, Definition 2.2 (B1)], a necessary condition for a map $\mu : M \rightarrow G$ to be a group valued moment map is $\mu^*\chi$ being exact, where χ is the canonical closed bi-invariant 3-form on G .

Let $\tilde{G}$ be as above, and define $\tilde{\mu} : \tilde{G}^{2l+1} \rightarrow \tilde{G}$ by

$$(a_1, b_1, \dots, a_l, b_l, c) \mapsto a_1b_1a_1^{-1}b_1^{-1} \cdots a_lb_la_l^{-1}b_l^{-1}c^2.$$

Let $Q_{2l+1} : \tilde{G}^{2l+1} \rightarrow \tilde{G}$ be the projection to the last factor. We have $\tilde{\mu}^*\chi = 2Q_{2l+1}^*\chi$, which represents a nontrivial cohomology class. So $\tilde{\mu}$ cannot be a group valued moment map, and we cannot apply [AMM, Theorem 7.2] directly to $\tilde{\mu}$ to conclude that $\tilde{X}_k = \tilde{\mu}^{-1}(k)$ is connected.

4. CONNECTED SUM OF AN EVEN NUMBER OF COPIES OF THE REAL PROJECTIVE PLANE

In this section, Σ is a compact nonorientable surface which is homeomorphic to the connected sum of $2l + 2$ copies of RP^2 , or equivalently, the connected sum of a Riemann surface of genus l and a Klein bottle, where $l > 1$.

Let G be a connected, compact, semisimple Lie group, and let $\rho : \tilde{G} \rightarrow G$ be the universal covering map. The space $\text{Hom}(\pi_1(\Sigma), G)$ can be identified with

$$X = \{(a_1, b_1, \dots, a_l, b_l, c_1, c_2) \in G^{2l+2} \mid a_1b_1a_1^{-1}b_1^{-1} \cdots a_lb_la_l^{-1}b_l^{-1}c_1^2c_2^2 = e\},$$

where e is the identity element of G .

Let $K = \text{Ker } \rho \cong \pi_1(G)$. There is an obstruction map

$$o_2 : \text{Hom}(\pi_1(\Sigma), G) \cong X \rightarrow K/2K$$

defined by

$$(a_1, b_1, \dots, a_l, b_l, c_1, c_2) \in X \mapsto \tilde{a}_1\tilde{b}_1\tilde{a}_1^{-1}\tilde{b}_1^{-1} \cdots \tilde{a}_l\tilde{b}_l\tilde{a}_l^{-1}\tilde{b}_l^{-1}\tilde{c}_1^2\tilde{c}_2^2 \in K/2K,$$

where $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l, \tilde{c}_1, \tilde{c}_2)$ is a preimage of $(a_1, b_1, \dots, a_l, b_l, c_1, c_2)$ under $\rho^{2l+2} : \tilde{G}^{2l+2} \rightarrow G^{2l+2}$. It is easily checked that the definition does not depend on the choice of $(\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_l, \tilde{b}_l, \tilde{c}_1, \tilde{c}_2)$ and o_2 descends to a continuous map

$$\bar{o}_2 : \text{Hom}(\pi_1(\Sigma), G)/G \rightarrow K/2K.$$

For each $k \in K$, define

$$\tilde{X}_k = \{(a_1, b_1, \dots, a_l, b_l, c_1, c_2) \in \tilde{G}^{2l+2} \mid a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1} c_1^2 c_2^2 = k\}.$$

Let $\bar{k}$ be the image of k under $K \rightarrow K/2K$. Then there is a surjective continuous map

$$\tilde{X}_k \rightarrow \bar{o}_2^{-1}(\bar{k})$$

which is the restriction of $\tilde{G}^{2l+2} \xrightarrow{\rho^{2l+2}} G^{2l+2} \rightarrow G^{2l+2}/G$, where G acts on $\tilde{G}^{2l+2}$ by diagonal conjugation. For each $k \in K$, $\tilde{X}_k$ contains $(\mu_{\tilde{G}}^l)^{-1}(k) \times \{e\} \times \{e\}$ which is nonempty by Fact 3. Therefore, $\bar{o}_2$ is surjective. It remains to show that $\tilde{X}_k$ is connected for all $k \in K$, which will imply that $\bar{o}_2^{-1}(\bar{k})$ is connected for all $\bar{k} \in K/2K$.

Let $Q' : \tilde{G}^{2l+2} \rightarrow \tilde{G}^2$ be the projection to the last two factors. The restriction of Q' to $\tilde{X}_k$ gives a map $Q : \tilde{X}_k \rightarrow \tilde{G}^2$, $(a_1, b_1, \dots, a_l, b_l, c_1, c_2) \mapsto (c_1, c_2)$. Fact 3 implies that Q is surjective and $Q^{-1}(c_1, c_2)$ is connected for all $(c_1, c_2) \in \tilde{G}^2$. Let $q \in T$ be a square root of $k \in Z(\tilde{G})$, where we fix a maximal torus T of $\tilde{G}$. To prove that $\tilde{X}_k$ is connected, it suffices to show that for any $(c_1, c_2) \in \tilde{G}^2$, there is a path $\gamma : [0, 1] \rightarrow \tilde{X}_k$ such that $\gamma(0) \in Q^{-1}(\tilde{e}, q)$ and $\gamma(1) \in Q^{-1}(c_1, c_2)$, where $\tilde{e}$ is the identity element of $\tilde{G}$.

For any $(c_1, c_2) \in \tilde{G}^2$, there exists $g_1, g_2 \in \tilde{G}$ such that $g_j^{-1} c_j g_j \in T$ for $j = 1, 2$. Let $\tilde{\mathfrak{g}}$ and $\mathfrak{t}$ be the Lie algebras of $\tilde{G}$ and T , respectively. Let $\exp : \tilde{\mathfrak{g}} \rightarrow \tilde{G}$ be the exponential map. Then $g_1^{-1} c_1 g_1 = \exp \xi_1$ for some $\xi_1 \in \mathfrak{t}$, and $g_2^{-1} c_2 g_2 q^{-1} = \exp \xi_2$ for some $\xi_2 \in \mathfrak{t}$. Let W be the Weyl group of $\tilde{G}$, and let $w \in W$ be a Coxeter element. The linear map $w : \mathfrak{t} \rightarrow \mathfrak{t}$ has no eigenvalue equal to 1, so there exists $\xi'_j \in \mathfrak{t}$ such that $w \cdot \xi'_j - \xi'_j = \xi_j$ for $j = 1, 2$, where $w = aT \in N(T)/T = W$ for some $a \in \tilde{G}$. We have

$$a \exp(t\xi'_j) a^{-1} \exp(-t\xi'_j) = \exp(t\xi_j)$$

for any $t \in \mathbb{R}$, where $j = 1, 2$.

The group $\tilde{G}$ is connected, so there exist paths $\tilde{g}_j : [0, 1] \rightarrow \tilde{G}$ such that $\tilde{g}_j(0) = \tilde{e}$ and $\tilde{g}_j(1) = g_j$ for $j = 1, 2$. Define $\gamma : [0, 1] \rightarrow \tilde{G}^{2l+2}$ by

$$\gamma(t) = (a_2(t), b_2(t), a_1(t), b_1(t), \tilde{e}, \dots, \tilde{e}, c_1(t), c_2(t))$$

where

$$a_j(t) = \tilde{g}_j(t) a \tilde{g}_j(t)^{-1}, \quad b_j(t) = \tilde{g}_j(t) \exp(-2t\xi'_j) \tilde{g}_j(t)^{-1}$$

for $j = 1, 2$, and

$$c_1(t) = \tilde{g}_1(t) \exp(t\xi_1) \tilde{g}_1(t)^{-1}, \quad c_2(t) = \tilde{g}_2(t) q \exp(t\xi_2) \tilde{g}_2(t)^{-1}.$$

Then the image of γ lies in $\tilde{X}_k$, $\gamma(0) = (a, \tilde{e}, a, \tilde{e}, \tilde{e}, \dots, \tilde{e}, q) \in Q^{-1}(\tilde{e}, q)$, and

$$\gamma(1) = (g_2 a g_2^{-1}, g_2 \exp(-2\xi'_2) g_2^{-1}, g_1 a g_1^{-1}, g_1 \exp(-2\xi'_1) g_1^{-1}, \tilde{e}, \dots, \tilde{e}, c_1, c_2) \in Q^{-1}(c_1, c_2).$$

Remark 5. Let $\tilde{G}$ be as above, and define $\tilde{\mu} : \tilde{G}^{2l+2} \rightarrow \tilde{G}$ by

$$(a_1, b_1, \dots, a_l, b_l, c_1, c_2) \mapsto a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_l b_l a_l^{-1} b_l^{-1} c_1^2 c_2^2.$$

Let Q_{2l+1} and Q_{2l+2} denote the projections from G^{2l+2} to the $(2l+1)$ -th and $(2l+2)$ -th factors, respectively. We have $\tilde{\mu}^* \chi = 2Q_{2l+1}^* \chi + 2Q_{2l+2}^* \chi$, which represents a nontrivial cohomology

class. So $\tilde{\mu}$ cannot be a group valued moment map, and we cannot apply [AMM, Theorem 7.2] directly to $\tilde{\mu}$ to conclude that $\tilde{X}_k = \tilde{\mu}^{-1}(k)$ is connected.

5. THE CASES $k = 1, 2, 4$

Let G be a compact, semisimple, connected and simply connected Lie group. By Theorem 2, $\text{Hom}(\pi_1(\Sigma), G)/G$ is connected if Σ is the connected sum of k copies of RP^2 , where $k \neq 1, 2, 4$. For $k = 1$, we have

$$\text{Hom}(\pi_1(RP^2), G)/G = \{g \in G \mid g^2 = e\}/G,$$

where G acts by conjugation. In particular, for $n > 1$,

$$\text{Hom}(\pi_1(RP^2), SU(n))/SU(n) \cong \left\{ \begin{pmatrix} -I_{2j} & 0 \\ 0 & I_{n-2j} \end{pmatrix} \mid j = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right\},$$

where I_m denotes the $m \times m$ identity matrix. For $n > 2$,

$$\text{Hom}(\pi_1(RP^2), Spin(n))/Spin(n) \cong \left\{ (-1)^j e_1 e_2 \cdots e_{2j} \mid j = 0, 1, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right\},$$

where we view $Spin(n)$ as a subset of the real Clifford algebra C_n , and $\{e_1, \dots, e_n\}$ is an orthonormal basis for $\mathbb{R}^n$. For $n > 0$,

$$\text{Hom}(\pi_1(RP^2), Sp(n))/Sp(n) \cong \left\{ \begin{pmatrix} -I_k & 0 & 0 \\ 0 & I_{n-k} & 0 \\ 0 & 0 & -I_k & 0 \\ & & 0 & I_{n-k} \end{pmatrix} \mid k = 0, 1, \dots, n \right\}.$$

So Theorem 2 is not true for $k = 1$.

It is not clear to us if Theorem 2 is true for $k = 2$. We expect Theorem 2 to hold for $k = 4$, but we do not know how to prove it using the approach in this paper.

APPENDIX A. OBSTRUCTION CLASS

Let Σ be a connected closed compact surface. Let G be a connected, compact, semisimple Lie group. Let $\rho : \tilde{G} \rightarrow G$ be the universal covering map which is also a group homomorphism. Then $\text{Ker} \rho$ is contained in the center $Z(\tilde{G})$ of $\tilde{G}$. Let $K = \text{Ker} \rho \cong \pi_1(G)$ which is a finite abelian group.

In this appendix, we will give several interpretations of the obstruction map

$$o_2 : \text{Hom}(\pi_1(\Sigma), G) \rightarrow H^2(\Sigma; K).$$

Note that the definition in Section 3 can be extended to the case $l = 0$, and the definition in Section 4 can be extended to the cases $l = 0, 1$. If Σ is homeomorphic to a 2-sphere, then $\text{Hom}(\pi_1(\Sigma), G)$ consists of a single point e . We extend the definition in Section 2 to the genus zero case by defining $o_2(e)$ to be the identity element of $H^2(\Sigma; K)$. So the obstruction map o_2 is defined for any closed compact surfaces Σ .

A.1. Non-abelian cohomology of groups. For an account of non-abelian cohomology of groups, see [Se, Chapter VII, Appendix].

We have a short exact sequence of groups

$$(2) \quad 1 \rightarrow K \rightarrow \tilde{G} \rightarrow G \rightarrow 1.$$

Let $\pi_1(\Sigma)$ act trivially on (2) on the left so that (2) can be viewed as a short exact sequence of (non-abelian) $\pi_1(\Sigma)$ -modules. The short exact sequence (2) gives rise to the following long exact sequence of pointed sets [Se, Chapter VII, Appendix, Proposition 2]:

$$(3) \quad \begin{array}{ccccccc} 1 & \rightarrow & H^0(\pi_1(\Sigma), K) & \rightarrow & H^0(\pi_1(\Sigma), \tilde{G}) & \rightarrow & H^0(\pi_1(\Sigma), G) \\ & & \xrightarrow{\delta} & & H^1(\pi_1(\Sigma), \tilde{G}) & \rightarrow & H^1(\pi_1(\Sigma), G) \xrightarrow{\Delta} H^2(\pi_1(\Sigma), K) \end{array}$$

which can be rewritten as

$$(4) \quad \begin{array}{ccccccc} 1 & \rightarrow & K & \rightarrow & \tilde{G} & \rightarrow & G \\ & & \xrightarrow{\delta} & & \text{Hom}(\pi_1(\Sigma), \tilde{G})/\tilde{G} & \rightarrow & \text{Hom}(\pi_1(\Sigma), G)/G \xrightarrow{\Delta} H^2(\pi_1(\Sigma), K) \end{array}$$

by checking the definitions.

Let $p \in \text{Hom}(\pi_1(\Sigma), G)$ represent a point $\bar{p} \in H^1(\pi_1(\Sigma), G) \cong \text{Hom}(\pi_1(\Sigma), G)/G$. From the exact sequence of (4), p can be lifted to a homomorphism $\tilde{p} : \pi_1(\Sigma) \rightarrow \tilde{G}$ if and only if $\Delta(\bar{p})$ is the identity element of $H^2(\pi_1(\Sigma), K)$. From the explicit expression of the obstruction map $o_2 : \text{Hom}(\pi_1(\Sigma), G) \rightarrow H^2(\Sigma; K)$ we see that $o_2(p)$ is the identity element of the group $H^2(\Sigma; K)$ if and only if p can be lifted to a homomorphism $\tilde{p} : \pi_1(\Sigma) \rightarrow \tilde{G}$. Actually,

$$\Delta : H^1(\pi_1(\Sigma), G) \rightarrow H^2(\pi_1(\Sigma), K)$$

coincides with

$$\bar{o}_2 : \text{Hom}(\pi_1(\Sigma), G)/G \rightarrow H^2(\Sigma; K)$$

under the identifications

$$H^1(\pi_1(\Sigma), G) \cong \text{Hom}(\pi_1(\Sigma), G)/G, \quad H^2(\pi_1(\Sigma), K) \cong H^2(\Sigma; K),$$

where $H^2(\pi_1(\Sigma), K) \cong H^2(\Sigma; K)$ because Σ is the Eilenberg-MacLane space $K(\pi_1(\Sigma), 1)$.

A.2. Non-abelian Čech cohomology. For an account of non-abelian Čech cohomology, see [LM, Appendix A].

A.2.1. Flat bundles. Given a group A , let A also denote the sheaf on Σ for which $A(U)$ is the group of locally constant functions from U to A , where U is any open subset of Σ . The short exact sequence of sheaves

$$1 \rightarrow K \rightarrow \tilde{G} \rightarrow G \rightarrow 1$$

gives rise to the following long exact sequences of non-abelian Čech cohomology [LM, Remark A.2]

$$(5) \quad \begin{array}{ccccccc} 1 & \rightarrow & H^0(\Sigma; K) & \rightarrow & H^0(\Sigma; \tilde{G}) & \rightarrow & H^0(\Sigma; G) \\ & & \xrightarrow{\delta} & & H^1(\Sigma; \tilde{G}) & \rightarrow & H^1(\Sigma; G) \xrightarrow{\Delta} H^2(\Sigma; K) \end{array}$$

which can be rewritten as

$$(6) \quad \begin{array}{ccccccc} 1 & \rightarrow & K & \rightarrow & \tilde{G} & \rightarrow & G \\ & \searrow \delta & \rightarrow H^1(\Sigma; K) & \rightarrow & H^1(\Sigma; \tilde{G}) & \rightarrow & H^1(\Sigma; G) \xrightarrow{\Delta} H^2(\Sigma; K) \end{array}$$

by checking the definitions. The pointed sets $H^1(\Sigma; \tilde{G})$ and $H^1(\Sigma; G)$ can be identified with the moduli spaces of gauge equivalence classes of flat $\tilde{G}$ -bundles and G -bundles, respectively. The map δ in (6) is the trivial group homomorphism, while the map Δ in (6) sends a gauge equivalence class of flat G -bundles to the obstruction to lifting it to a gauge equivalence class of flat $\tilde{G}$ -bundles. Actually,

$$\Delta : H^1(\Sigma; G) \rightarrow H^2(\Sigma; K)$$

coincides with

$$\bar{o}_2 : \text{Hom}(\pi_1(\Sigma), G)/G \rightarrow H^2(\Sigma; K)$$

under the identification $H^1(\Sigma; G) \cong \text{Hom}(\pi_1(\Sigma), G)/G$.

A.2.2. Topological bundles. Given a group A , let $\underline{A}$ denote the sheaf on Σ for which $\underline{A}(U)$ is the group of continuous functions from U to A , where U is any open subset of Σ . Note that $\underline{K} = K$ since K is discrete. The short exact sequence of sheaves

$$1 \rightarrow K \rightarrow \underline{\tilde{G}} \rightarrow \underline{G} \rightarrow 1$$

gives rise to the following long exact sequence of Čech cohomology

$$(7) \quad \begin{array}{ccccccc} 1 & \rightarrow & H^0(\Sigma; K) & \rightarrow & H^0(\Sigma; \underline{\tilde{G}}) & \rightarrow & H^0(\Sigma, \underline{G}) \\ & \searrow \delta' & \rightarrow H^1(\Sigma; K) & \rightarrow & H^1(\Sigma; \underline{\tilde{G}}) & \rightarrow & H^1(\Sigma; \underline{G}) \xrightarrow{\Delta'} H^2(\Sigma; K). \end{array}$$

The pointed sets $H^1(\Sigma; \underline{\tilde{G}})$ and $H^1(\Sigma; \underline{G})$ can be identified with the set of equivalence classes of topological principal $\tilde{G}$ -bundles and G -bundles, respectively. Note that $H^1(\Sigma; K) \cong \text{Hom}(\pi_1(\Sigma), K)$. The exact sequence (7) can be rewritten as

$$(8) \quad \begin{array}{ccccccc} 1 & \rightarrow & K & \rightarrow & \text{Map}(\Sigma, \tilde{G}) & \rightarrow & \text{Map}(\Sigma, G) \\ & \searrow \delta' & \rightarrow \text{Hom}(\pi_1(\Sigma), K) & \rightarrow & \text{Prin}_{\tilde{G}}(\Sigma) & \rightarrow & \text{Prin}_G(\Sigma) \xrightarrow{\Delta'} H^2(\Sigma, K). \end{array}$$

The map δ' sends $f : \Sigma \rightarrow G$ to $f_* : \pi_1(\Sigma) \rightarrow \pi_1(G) \cong K$, and Δ' sends a principal G -bundle to the obstruction to lifting it to a principal $\tilde{G}$ -bundle. For example, if $G = SO(n)$ ($n > 2$), then $K = \mathbb{Z}/2\mathbb{Z}$, and $\Delta'(P)$ is the second Stiefel-Whitney class $w_2(P) \in H^2(\Sigma, \mathbb{Z}/2\mathbb{Z})$.

We have a commutative diagram of sheaves

$$\begin{array}{ccccccccc} 1 & \longrightarrow & K & \longrightarrow & \tilde{G} & \longrightarrow & G & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & K & \longrightarrow & \underline{\tilde{G}} & \longrightarrow & \underline{G} & \longrightarrow & 1 \end{array}$$

which induces a morphism from (5) to (7). In particular, we have the following commutative diagram

$$\begin{array}{ccc} H^1(\Sigma; G) = \text{Hom}(\pi_1(\Sigma), G)/G & \xrightarrow{\Delta} & H^2(\Sigma; K) \\ i \downarrow & & id \downarrow \\ H^1(\Sigma; \underline{G}) = \text{Prin}_G(\Sigma) & \xrightarrow{\Delta'} & H^2(\Sigma; K) \end{array}$$

where i sends a flat G -bundle to the underlying topological principal G -bundle. Therefore, if P is the underlying topological principal G -bundle of a flat G -bundle associated to $p \in \text{Hom}(\pi_1(\Sigma), G)$, then $\Delta'(P) = o_2(p) \in H^2(\Sigma; K)$.

A.3. Obstruction theory. For an account of obstruction theory of fiber bundles, see [St].

The first obstruction to the triviality of a principal G -bundle P over Σ , or equivalently, the first obstruction of existence of a cross-section, lies in $H^1(\Sigma; \pi_0(G))$, which vanishes because G is connected. So P is trivial over the 1-skeleton. The second obstruction lies in $H^2(\Sigma; \pi_1(G)) = H^2(\Sigma; K)$ and coincides with $\Delta'(P)$, the obstruction to lifting P to a principal $\tilde{G}$ -bundle. For a surface Σ , $\Delta'(P)$ is the only obstruction to the triviality of P . So the topological principal G -bundles over Σ are classified by $\Delta'(P) \in H^2(\Sigma; K)$. The set $\text{Prin}_{\tilde{G}}(\Sigma)$ consists of a single point corresponding to the trivial $\tilde{G}$ -bundle, and Δ' in (7) is a bijection.

Note that if G is not simply connected, then there exists nontrivial topological G -bundles over S^2 , but the underlying topological principal G -bundle of a flat G -bundle over S^2 must be trivial. So Δ sends the only point in $H^1(S^2; G) = H^1(\pi_1(S^2), G)$ to the identity element of $H^2(S^2; K) = H^2(\pi_1(S^2), K)$.

A.4. Elementary approach. There is an elementary approach to the classification of topological principal G -bundles over Σ which we learned from E. Meinrenken. Let D be a disc around a point $x \in \Sigma$. Any principal G -bundle P is trivial over D and over $\Sigma \setminus \{x\}$, and the topological type of P is determined by the homotopy class of the transition function $\psi : D \setminus \{x\} \rightarrow G$, or equivalently, an element in $K = \pi_1(G)$. This gives a surjective map $K \rightarrow \text{Prin}_G(\Sigma)$ which is injective if Σ is orientable and induces a bijection $K/2K \rightarrow \text{Prin}_G(\Sigma)$ if Σ is nonorientable. Therefore, we have a bijection

$$\phi : H^2(\Sigma; K) \rightarrow \text{Prin}_G(\Sigma).$$

The argument in [Li, Section 6] shows that if P is the underlying topological principal G -bundle of a flat G -bundle associated to $p \in \text{Hom}(\pi_1(\Sigma), G)$, then $\phi(o_2(p)) = P$.

Based on above discussions, Theorem 1 and Theorem 2 can be reformulated as follows.

Theorem 6. *Let Σ be a closed compact orientable surface of genus $l > 0$, or a closed compact nonorientable surface which is homeomorphic to k copies of the real projective plane, where $k \neq 1, 2, 4$. Let G be a connected, compact, semisimple Lie group, and let P be a topological principal G -bundle over Σ . Then the moduli space of gauge equivalence classes of flat connections on P is nonempty and connected.*

REFERENCES

- [AB] M.F. Atiyah and R. Bott, *The Yang-Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A **308** (1983), no. 1505, 523–615.
- [AMM] A. Alekseev, A. Malkin, and E. Meinrenken, *Lie group valued moment maps*, J. Differential Geom. **48** (1998), no. 3, 445–495.
- [AMW] A. Alekseev, E. Meinrenken, and C. Woodward, *Duistermaat-Heckman measures and moduli spaces of flat bundles over surfaces*, Geom. Funct. Anal. **12** (2002), no. 1, 1–31.
- [BD] Theodor Bröcker and Tammo tom Dieck, *Representations of compact Lie groups*, Graduate Texts in Mathematics, 98. Springer-Verlag, New York, 1985.
- [Go] W.M. Goldman, *Topological components of spaces of representations*, Invent. Math. **93** (1988), no. 3, 557–607.
- [GM] W.M. Goldman and J.J. Millson, *The deformation theory of representations of fundamental groups of compact Kähler manifolds*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), no. 2, 153–158.
- [HM] K.H. Hofmann and S.A. Morris, *The structure of compact groups. A primer for the student—a handbook for the expert*, de Gruyter Studies in Mathematics, 25. Walter de Gruyter & Co., Berlin, 1998.
- [Hu] J.E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, 29. Cambridge University Press, Cambridge, 1990.
- [Li] J. Li, *The space of surface group representations*, Manuscripta Math. **78** (1993), no. 3, 223–243.
- [LM] H.B. Lawson and M. Michelsohn, *Spin geometry*, Princeton Mathematical Series 38, Princeton University Press, Princeton, NJ, 1989.
- [Mi] J. Milnor, *On the existence of a connection with curvature zero*, Comment. Math. Helv. **32** (1958), 215–223.
- [Ra] A. Ramanathan, *Moduli for principal bundles*, Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, Copenhagen, 1978), pp. 527–533, Lecture Notes in Math., 732, Springer, Berlin, 1979.
- [Se] J.-P. Serre, *Local fields*, translated from the French by Marvin Jay Greenberg, Graduate Texts in Mathematics, 67. Springer-Verlag, New York-Berlin, 1979.
- [St] N. Steenrod, *The Topology of Fibre Bundles*, Princeton Mathematical Series, vol. 14. Princeton University Press, Princeton, N. J., 1951.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO

E-mail address: `nankuo@math.toronto.edu`

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY

E-mail address: `ccliu@math.harvard.edu`